

## MODIFIED RANKIN SCALE (MRS) for children

Date: \_\_\_\_\_

Rater Name: \_\_\_\_\_

Score	Description
0	No symptoms at all
1	No significant disabilities despite symptoms in clinical examination; age appropriate behaviour and further development
2	Slight disability; unable to carry out all previous activities, but same independence as other age- and sex-matched children (no reduction of levels on the gross motor function scale )
3	Moderate disability; requiring some help, but able to walk without assistance; in younger patients adequate motor development despite mild functional impairment (reduction of one level on the gross motor function scale)
4	Moderately severe disability; unable to walk without assistance; in younger patients reduction of at least 2 levels on the gross motor function scale
5	Severe disability; bedridden, requiring constant nursing care and attention
6	Dead

Score (0 – 6): \_\_\_\_\_

### References

Rankin J. "Cerebral vascular accidents in patients over the age of 60." *Scott Med J* 1957;2:200-15

Bonita R, Beaglehole R. "Modification of Rankin Scale: Recovery of motor function after stroke." *Stroke* 1988 Dec;19(12):1497-1500

Van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. "Interobserver agreement for the assessment of handicap in stroke patients." *Stroke* 1988;19(5):604-7

Palisano R, Rosenbaum P, Walter S, Russel D, Wood E, Galuppi B. "Development and reliability of a system to classify gross motor function in children with cerebral palsy" *Dev Med Child Neurol.* 1997;39:214e23.

Bigi S, Fischer U et al. "Acute ischaemic stroke in children and young adults: Differences in clinical presentation, aetiology and outcome" *Ann Neurol (in revision)*